

On The Location of Zeros of Polynomials with Different Complex Coefficients

P.Ramulu^{1‡}, C.Gangadhar¹, M.Amarnath² and G.L.Reddy³

^{1‡}*Department of Mathematics, M.V.S. Govt. Arts and Science College(A),
.....Mahabubnagar, Telangana-509001, India.*

²*Department of Mathematics, National Institute of Technology, Andhra Pradesh, India.*

²*Department of Mathematics, CBIT, Hyderabad, Telangana.*

³*School of Mathematics and Statistics, University of Hyderabad, India 500046. ,*

ABSTRACT

In this paper we extend Eneström-Keakeya theorem (Let $P(z) = \sum_{i=0}^n a_i z^i$ be a polynomial of degree n such that $0 < a_0 \leq a_1 \leq \dots \leq a_n$, then all the zeros of $P(z)$ lie in $|z| \leq 1$) for polynomials with complex coefficients the result of [8] have been generalized by relaxing the hypothesis in different ways by considering complex coefficients.

Mathematics Subject Classification: 30C10, 30C15.[2pt]

Keywords: Zeros of polynomial, Eneström-Keakeya theorem.

1. INTRODUCTION

Location of zeros of a polynomial is a long standing classical problem [1,3-5,8,9,11-13]. It is an interesting area of research for engineers as well as mathematicians and many results on the same topic are available in literature. Here we make an attempt to extend some of the known result for real coefficients to complex coefficients. Existing results in the literature also show that there is a need to find bounds for special polynomials, for example, for those having restrictions on the coefficients, there is always a need for refinement of results in this subject. The well known result in the theory of distribution of zeros of polynomials is the following:

Theorem A₁. [2, 7] : Let $P(z) = \sum_{i=0}^n a_i z^i$ be a polynomial of degree n such that $0 < a_0 \leq a_1 \leq \dots \leq a_{n-1} \leq a_n$ then all the zeros of $P(z)$ lie in $|z| \leq 1$.

A.Joyal, G.Labelle and Q. I. Rahman [6] obtained the following generalization, by considering the coefficients to be real, instead of being only positive.

Theorem A₂. Let $P(z) = \sum_{i=0}^n a_i z^i$ be a polynomial of degree n such that $a_0 \leq a_1 \leq \dots \leq a_{n-1} \leq a_n$ then all the zeros of $P(z)$ lie in $|z| \leq \frac{1}{|a_n|} \{a_n - a_0 + |a_0|\}$.

Theorem A₃. [10] : Let $P(z) = \sum_{i=0}^n a_i z^i$ be a polynomial of degree n with real coefficients such that

$$a_0 \leq a_1 \leq \dots \leq a_{m-1} \leq a_m \geq a_{m+1} \geq \dots \geq a_{n-1} \geq a_n.$$

Then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|a_n|} [2a_m + |a_0| - (a_0 + |a_n|)].$$

Theorem A₄. [8] : Let $P(z) = \sum_{i=0}^n a_i z^i$ be a polynomial of degree $n \geq 2$ with real coefficients such that $a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0$ if n is even

OR

$a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0$ if n is odd

then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|a_n|} \{ |a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] - [a_{n-1} + a_{n-3} + \dots + a_3 + a_1]) \}$$

if n is even (OR)

$$|z| \leq \frac{1}{|a_n|} \{ a_n + 2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1] - [a_{n-1} + a_{n-3} + \dots + a_4 + a_2]) \}$$

if n is odd

In this paper we want to prove the following results.

Theorem 1. Let $P(z) = \sum_{j=0}^n \alpha_j z^j$ be a polynomial with complex coefficients of degree $n \geq 2$ with $Re(\alpha_j) = a_j$ and $Im(\alpha_j) = b_j$ for $j = 0, 1, 2, \dots, n$ such that

$a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0$ and

$b_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \geq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0$ if n is even

OR

$a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0$ and

$b_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \geq \dots \geq b_4 \leq b_3 \leq b_2 \leq b_1 \geq b_0$ if n is odd

then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} \left[|a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) + |b_0| + b_0 + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2] - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \right] \text{ if } n \text{ is even}$$

OR

$$|z| \leq \frac{1}{|\alpha_n|} \left[|a_0| - a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1] - [a_{n-1} + a_{n-3} + \dots a_4 + a_2]) + |b_0| - b_0 + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_3 + b_1] - [b_{n-1} + b_{n-3} \dots + b_4 + b_2]) \right] \text{ if } n \text{ is odd}$$

Corollary 1. Let $P(z) = \sum_{j=0}^n \alpha_j z^j$ be a polynomial with complex coefficients of degree $n \geq 2$ with $Re(\alpha_j) = a_j \geq 0$ and $Im(\alpha_j) = b_j \geq 0$ for $j = 0, 1, 2, \dots, n$ such that

$$a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0 \text{ and}$$

$$b_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \geq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0 \text{ if } n \text{ is even}$$

OR

$$a_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \geq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0 \text{ and}$$

$$b_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \geq \dots \geq b_4 \leq b_3 \leq b_2 \leq b_1 \geq b_0 \text{ if } n \text{ is odd}$$

then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} \left[a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2 + a_0] - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2 + b_0] - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \right] \text{ if } n \text{ is even}$$

OR

$$|z| \leq \frac{1}{|\alpha_n|} \left[a_n + 2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1] - [a_{n-1} + a_{n-3} + \dots a_4 + a_2]) + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_3 + b_1] - [b_{n-1} + b_{n-3} \dots + b_4 + b_2]) \right] \text{ if } n \text{ is odd}$$

Remark 1. By taking $a_j \geq 0, b_j \geq 0$ for $j = 0, 1, 2, \dots, n$ in the Theorem 1, then it reduces to Corollary 1

Theorem 2. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial with complex coefficients of degree $n \geq 2$ with $Re(\alpha_i) = a_i$ and $Im(\alpha_i) = b_i$ for $i = 0, 1, 2, 3, \dots, n$ such that for some $k \geq 1, \delta \geq 0,$

$ka_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0 + \delta$ and
 $kb_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0 + \delta$ if n is even

OR

$ka_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0 - \delta$ and
 $kb_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0 - \delta$ if n is odd

then all the zeros of the polynomial $P(z)$ lie in

$$|z+k-1| \leq \frac{1}{|\alpha_n|} \{ka_n + |a_0| + a_0 + 2\delta + 2[(a_{n-2} + a_{n-4} + \dots + a_4 + a_2) - (a_{n-1} + a_{n-3} + \dots + a_3 + a_1)] + kb_n + |b_0| + b_0 + 2\delta + 2[(b_{n-2} + b_{n-4} + \dots + b_4 + b_2) - (b_{n-1} + b_{n-3} + \dots + b_3 + b_1)]\}$$

if n is even

OR

$$|z+k-1| \leq \frac{1}{|\alpha_n|} \{ka_n + |a_0| - a_0 + 2\delta + 2[(a_{n-2} + a_{n-4} + \dots + a_3 + a_1) - (a_{n-1} + a_{n-3} + \dots + a_4 + a_2)] + kb_n + |b_0| - b_0 + 2\delta + 2[(b_{n-2} + b_{n-4} + \dots + b_3 + b_1) - (b_{n-1} + b_{n-3} + \dots + b_4 + b_2)]\}$$

if n is odd

Corollary 2. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial with complex coefficients of degree $n \geq 2$ with $Re(\alpha_i) = a_i$ and $Im(\alpha_i) = b_i$ for $i = 0, 1, 2, 3, \dots, n$ such that for some $k \geq 1$,

$ka_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0$ and
 $kb_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0$ if n is even

OR

$ka_n \geq a_{n-1} \leq a_{n-2} \geq a_{n-3} \leq a_{n-4} \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0$ and
 $kb_n \geq b_{n-1} \leq b_{n-2} \geq b_{n-3} \leq b_{n-4} \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0$ if n is odd

then all the zeros of the polynomial $P(z)$ lie in

$$|z+k-1| \leq \frac{1}{|\alpha_n|} \{ka_n + |a_0| + a_0 + 2[(a_{n-2} + a_{n-4} + \dots + a_4 + a_2) - (a_{n-1} + a_{n-3} + \dots + a_3 + a_1)] + kb_n + |b_0| + b_0 + 2[(b_{n-2} + b_{n-4} + \dots + b_4 + b_2) - (b_{n-1} + b_{n-3} + \dots + b_3 + b_1)]\}$$

if n is even

OR

$$|z+k-1| \leq \frac{1}{|\alpha_n|} \{ka_n + |a_0| - a_0 + 2[(a_{n-2} + a_{n-4} + \dots + a_3 + a_1) - (a_{n-1} + a_{n-3} + \dots + a_4 + a_2)] + kb_n + |b_0| - b_0 + 2[(b_{n-2} + b_{n-4} + \dots + b_3 + b_1) - (b_{n-1} + b_{n-3} + \dots + b_4 + b_2)]\}$$

if n is odd

Remark 2. By taking $\delta = 0$ in the Theorem 2, then it reduces to Corollary 2

Theorem 3. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial with complex coefficients degree $n \geq 2$ with $Re(\alpha_i) = a_i$ and $Im(\alpha_i) = b_i$ for $i = 0, 1, 2, \dots, n$ such that

$a_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0$,
 $b_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0$ if n is even

OR

$a_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0$,
 $b_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0$ if n is odd

then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} [|a_0| - a_0 - a_n + 2([a_{n-1} + a_{n-3} + \dots + a_3 + a_1] - [a_{n-2} + a_{n-4} + \dots + a_4 + a_2]) + |b_0| - b_0 - b_n + 2([b_{n-1} + b_{n-3} + \dots + b_3 + b_1] - [b_{n-2} + b_{n-4} + \dots + b_4 + b_2])]$$

if n is even

OR

$$|z| \leq \frac{1}{|\alpha_n|} [|a_0| + a_0 - a_n + 2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1] - [a_{n-1} + a_{n-3} + \dots + a_4 + a_2]) + |b_0| + b_0 - b_n + 2([b_{n-2} + b_{n-4} + \dots + b_3 + b_1] - [b_{n-1} + b_{n-3} + \dots + b_4 + b_2])]$$

if n is odd

Corollary 3. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial with complex coefficients degree $n \geq 2$ with $Re(\alpha_i) = a_i \geq 0$ and $Im(\alpha_i) = b_i \geq 0$ for $i = 0, 1, 2, \dots, n$ such that

$$a_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0,$$

$$b_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0 \text{ if } n \text{ is even}$$

OR

$$a_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0,$$

$$b_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0 \text{ if } n \text{ is odd}$$

then all the zeros of $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} [2([a_{n-1} + a_{n-3} + \dots + a_3 + a_1] - [a_{n-2} + a_{n-4} + \dots + a_4 + a_2]) - a_n + 2([b_{n-1} + b_{n-3} + \dots + b_3 + b_1] - [b_{n-2} + b_{n-4} + \dots + b_4 + b_2]) - b_n] \text{ if } n \text{ is even}$$

OR

$$|z| \leq \frac{1}{|\alpha_n|} [2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1 + a_0] - [a_{n-1} + a_{n-3} + \dots + a_4 + a_2]) - a_n + 2([b_{n-2} + b_{n-4} + \dots + b_3 + b_1 + b_0] - [b_{n-1} + b_{n-3} + \dots + b_4 + b_2]) - b_n] \text{ if } n \text{ is odd}$$

Remark 3. By taking $Re(\alpha_i) = a_i \geq 0, Im(\alpha_i) = b_i \geq 0$ for $i = 0, 1, 2, \dots, n$ in the Theorem 3, then it reduces to Corollary 3.

Theorem 4. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial with complex coefficients of degree $n \geq 2$ with $Re(\alpha_i) = a_i$ and $Im(\alpha_i) = b_i$ for $i = 0, 1, 2, \dots, n$ such that for $0 \leq r, s \leq 1, \delta > 0, \eta > 0$

$$ra_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0 - \delta,$$

$$sb_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0 - \eta \text{ if } n \text{ is even}$$

OR

$$ra_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0 + \delta,$$

$$sb_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0 + \eta \text{ if } n \text{ is odd}$$

then all the zeros of the polynomial $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} \{ 2\delta + |a_n| + |a_0| - a_0 - r(|a_n| + a_n) + 2([a_{n-1} + a_{n-3} + \dots + a_3 + a_1] - [a_{n-2} + a_{n-4} + \dots + a_4 + a_2]) + 2\eta + |b_n| + |b_0| - b_0 - s(|b_n| + b_n) + 2([b_{n-1} + b_{n-3} + \dots + b_3 + b_1] - [b_{n-2} + b_{n-4} + \dots + b_4 + b_2]) \} \text{ if } n \text{ is even}$$

OR

$$|z| \leq \frac{1}{|\alpha_n|} \{2\delta + |a_n| + |a_0| + a_0 - r(|a_n| + a_n) + 2[(a_{n-1} + a_{n-3} + \dots + a_4 + a_2) - (a_{n-2} + a_{n-4} + \dots + a_3 + a_1)] + 2\eta + |b_n| + |b_0| + b_0 - s(|b_n| + b_n) + 2[(b_{n-1} + b_{n-3} + \dots + b_4 + b_2) - (b_{n-2} + b_{n-4} + \dots + b_3 + b_1)]\} \text{ if } n \text{ is odd}$$

Corollary 4. Let $P(z) = \sum_{i=0}^n \alpha_i z^i$ be a polynomial of degree ≥ 2 with $Re(\alpha_i) = a_i$ and $Im(\alpha_i) = b_i$ for $i = 0, 1, 2, \dots, n$ such that for $0 \leq r, s \leq 1$

$$ra_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \geq a_4 \leq a_3 \geq a_2 \leq a_1 \geq a_0,$$

$$sb_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \geq b_4 \leq b_3 \geq b_2 \leq b_1 \geq b_0 \text{ if } n \text{ is even}$$

OR

$$ra_n \leq a_{n-1} \geq a_{n-2} \leq a_{n-3} \geq a_{n-4} \leq \dots \leq a_4 \geq a_3 \leq a_2 \geq a_1 \leq a_0,$$

$$sb_n \leq b_{n-1} \geq b_{n-2} \leq b_{n-3} \geq b_{n-4} \leq \dots \leq b_4 \geq b_3 \leq b_2 \geq b_1 \leq b_0 \text{ if } n \text{ is odd}$$

then all the zeros of the polynomial $P(z)$ lie in

$$|z| \leq \frac{1}{|\alpha_n|} \{|a_n| + |a_0| - a_0 - r(|a_n| + a_n) + 2[(a_{n-1} + a_{n-3} + \dots + a_3 + a_1) - (a_{n-2} + a_{n-4} + \dots + a_4 + a_2)] + |b_n| + |b_0| - b_0 - s(|b_n| + b_n) + 2[(b_{n-1} + b_{n-3} + \dots + b_3 + b_1) - (b_{n-2} + b_{n-4} + \dots + b_4 + b_2)]\} \text{ if } n \text{ is even}$$

OR

$$|z| \leq \frac{1}{|\alpha_n|} \{|a_n| + |a_0| + a_0 - r(|a_n| + a_n) + 2[(a_{n-1} + a_{n-3} + \dots + a_4 + a_2) - (a_{n-2} + a_{n-4} + \dots + a_3 + a_1)] + |b_n| + |b_0| + b_0 - s(|b_n| + b_n) + 2[(b_{n-1} + b_{n-3} + \dots + b_4 + b_2) - (b_{n-2} + b_{n-4} + \dots + b_3 + b_1)]\} \text{ if } n \text{ is odd}$$

Remark 4. By taking $\delta = 0$, $\eta = 0$ in the Theorem 4, then it reduces to Corollary 4.

2. PROOF OF THE THEOREMS

Proof of Theorem 1. Let $P(z) = \alpha_n z^n + \alpha_{n-1} z^{n-1} + \dots + \alpha_3 z^3 + \alpha_2 z^2 + \alpha_1 z + \alpha_0$ be a polynomial with complex of degree $n \geq 2$ with $\alpha_j = a_j + ib_j$ for $j = 0, 1, 2, \dots, n$.

Then consider the polynomial

$$\begin{aligned} Q(z) &= (1 - z)P(z) \\ &= -\alpha_n z^{n+1} + (\alpha_n - \alpha_{n-1})z^n + (\alpha_{n-1} - \alpha_{n-2})z^{n-1} \\ &\quad + (\alpha_{n-2} - \alpha_{n-3})z^{n-2} + \dots + (\alpha_1 - \alpha_0)z + \alpha_0. \\ &= -\alpha_n z^{n+1} + (a_n - a_{n-1})z^n + (a_{n-1} - a_{n-2})z^{n-1} \\ &\quad + (a_{n-2} - a_{n-3})z^{n-2} + \dots + (a_1 - a_0)z + a_0 \\ &\quad + i\{(b_n - b_{n-1})z^n + (b_{n-1} - b_{n-2})z^{n-1} \\ &\quad + (b_{n-2} - b_{n-3})z^{n-2} + \dots + (b_1 - b_0)z + b_0\}. \end{aligned}$$

If $|z| < 1$ then $\frac{1}{|z|^{n-i}} > 1$ for $i = 0, 1, 2, \dots, n - 1$.

$$\begin{aligned}
 \text{Now } |Q(z)| &\geq |\alpha_n||z|^{n+1} - \{|a_n - a_{n-1}||z|^n + |a_{n-1} - a_{n-2}||z|^{n-1} \\
 &\quad + |a_{n-2} - a_{n-3}||z|^{n-2} + \dots + |a_3 - a_2||z|^3 \\
 &\quad + |a_2 - a_1||z|^2 + |a_1 - a_0||z| + |a_0| \\
 &\quad + |b_n - b_{n-1}||z|^n + |b_{n-1} - b_{n-2}||z|^{n-1} \\
 &\quad + |b_{n-2} - b_{n-3}||z|^{n-2} + \dots \\
 &\quad + |b_3 - b_2||z|^3 + |b_2 - b_1||z|^2 + |b_1 - b_0||z| + |b_0|\} \\
 &\geq |\alpha_n||z|^n \left[|z| - \frac{1}{|\alpha_n|} \{|a_n - a_{n-1}| \right. \\
 &\quad + \frac{|a_{n-1} - a_{n-2}|}{|z|} + \frac{|a_{n-2} - a_{n-3}|}{|z|^2} + \dots + \frac{|a_3 - a_2|}{|z|^{n-3}} \\
 &\quad + \frac{|a_2 - a_1|}{|z|^{n-2}} + \frac{|a_1 - a_0|}{|z|^{n-1}} + \frac{|a_0|}{|z|^n} + |b_n - b_{n-1}| \\
 &\quad + \frac{|b_{n-1} - b_{n-2}|}{|z|^1} + \frac{|b_{n-2} - b_{n-3}|}{|z|^2} + \dots \\
 &\quad \left. + \frac{|b_3 - b_2|}{|z|^{n-3}} + \frac{|b_2 - b_1|}{|z|^{n-2}} + \frac{|b_1 - b_0|}{|z|^n} + \frac{|b_0|}{|z|^n} \} \right] \\
 &\geq |\alpha_n||z|^n \left[|z| - \frac{1}{|\alpha_n|} \{|a_n - a_{n-1}| + |a_{n-1} - a_{n-2}| \right. \\
 &\quad + |a_{n-2} - a_{n-3}| + \dots + |a_3 - a_2| + |a_2 - a_1| + |a_1 - a_0| \\
 &\quad + |a_0| + |b_n - b_{n-1}| + |b_{n-1} - b_{n-2}| + |b_{n-2} - b_{n-3}| + \dots + |b_3 - b_2| \\
 &\quad \left. + |b_2 - b_1| + |b_1 - b_0| + |b_0|\} \right] \\
 &\geq |\alpha_n||z|^n \left[|z| - \frac{1}{|\alpha_n|} \{(a_n - a_{n-1}) \right. \\
 &\quad + (a_{n-1} - a_{n-2}) + (a_{n-2} - a_{n-3}) + \dots + (a_2 - a_3) + (a_2 - a_1) \\
 &\quad + (a_0 - a_1) + |a_0| + (b_n - b_{n-1}) + (b_{n-1} - b_{n-2}) \\
 &\quad + (b_{n-2} - b_{n-3}) + \dots + (b_2 - b_3) + (b_2 - b_1) \\
 &\quad \left. + (b_0 - b_1) + |b_0|\} \right] \text{ if } n \text{ is even, by hypothesis} \\
 &= |\alpha_n||z|^n \left[|z| - \frac{1}{|\alpha_n|} \{ \right. \\
 &\quad \{ |a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] \\
 &\quad - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) \\
 &\quad + |b_0| + b_0 + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2] \\
 &\quad - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \} \} \right] > 0
 \end{aligned}$$

$$\begin{aligned} \text{if } |z| > \{ & |a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] \\ & - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) + |b_0| + b_0 + b_n \\ & + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2] - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \} \end{aligned}$$

This shows that if $|z| > 1$ then $|Q(z)| > 0$ whenever

$$\begin{aligned} |z| > \frac{1}{|\alpha_n|} \{ & |a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] \\ & - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) + |b_0| + b_0 + b_n \\ & + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2] - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \} \end{aligned}$$

Hence all the zeros of $Q(z)$ with $|z| > 1$ lie in

$$\begin{aligned} |z| < \frac{1}{|\alpha_n|} \{ & |a_0| + a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_4 + a_2] \\ & - [a_{n-1} + a_{n-3} + \dots a_3 + a_1]) + |b_0| + b_0 + b_n \\ & + 2([b_{n-2} + b_{n-4} + \dots + b_4 + b_2] - [b_{n-1} + b_{n-3} \dots + b_3 + b_1]) \} \text{ if } n \text{ is even} \end{aligned}$$

But the zeros of $Q(z)$ whose modules is less than or equal to 1 already satisfy the above inequality. Since all the zeros of $Q(z)$ lie in the circle defined by the above inequality, we conclude that the proof of the Theorem 1 is complete, if n is even. Similarly we can also prove for odd degree polynomials. For this we can rearrange the terms of the given polynomial and compute as above. That is if n is odd then all the zeros of $P(z)$ lie in

$$\begin{aligned} |z| \leq \frac{1}{|\alpha_n|} \left[& |a_0| - a_0 + a_n + 2([a_{n-2} + a_{n-4} + \dots + a_3 + a_1] \right. \\ & - [a_{n-1} + a_{n-3} + \dots a_4 + a_2]) \\ & + |b_0| - b_0 + b_n + 2([b_{n-2} + b_{n-4} + \dots + b_3 + b_1] \\ & \left. - [b_{n-1} + b_{n-3} \dots + b_4 + b_2]) \right]. \end{aligned}$$

This completes the proof of the Theorem 1.

Proof of Theorem 2. Proof of theorem 2 is similar to the proof of theorem 1.

Proof of Theorem 3. Proof of theorem 3 is similar to the proof of theorem 1.

Proof of Theorem 4. Proof of theorem 4 is similar to the proof of theorem 1.

REFERENCES

- [1] A. Aziz and B. A. Zargar, Some extensions of Eneström-Kakeya Theorem, *Glasnik Matematički*, 31, (1996), 239-244.
- [2] G.Eneström, Remarque sur un théorème relatif aux racines de l'équation $a_n + \dots + a_0 = 0$ où tous les coefficients sont et positifs,
- [3] R. Fricke, *Lehrbuch der Algebra*, Friedrich Vieweg and Sohn Verlag, Braunschweig, Germany, 1924.
- [4] Gulshan Singh, "On the Eneström-Kakeya Theorem." *American Journal of Mathematical Analysis*, vol. 2, no. 1 (2014): 15-18.
- [5] W. Heitzinger, W. I. Troch and G. Valentin, *Praxis nichtlinearer Gleichungen*, Carl Hanser Verlag, München, Wien, Germany, Austria, 1985.
- [6] A. Joyal, G. Labelle and Q. I. Rahman, On the location of zeros of polynomials, *Canad. Math. Bull.* 10, (1967), 53-63.
- [7] S. Kakeya, On the limits of the roots of an algebraic equation with positive coefficients, *Tôhoku Math.J* 2 (1912-1913), 140-142.
- [8] P.Ramulu and G.L.Reddy, Location of zeros of polynomial with restricted real coefficients, *International J. of Math.Sci.&Engg. Appls(IJMSEA)*.ISSN0973-9424, Vol.9 No.II(June,2015),pp:1-12.
- [9] P. Ramulu and G.L. Reddy (2020). Location of Zeros of Polynomials, *Communications in Nonlinear Analysis*, 2020, 9(1).
- [10] P Ramulu, G.L.Reddy and C.Gangadhar, On the zeros of polynomials with real coefficients, *International Journal of Applied Mathematical Sciences*, 5(2), (2015), 109-118.
- [11] P.Ramulu and G.L.Reddy, On the Enestrom-Kakeya theorem, *International Journal of Pure and Applied Mathematics*, 102(4),(2015), 687-700.
- [12] W. M. Shah, A. Liman and Shamim Ahmad Bhat, On the Eneström-Kakeya Theorem, *International Journal of Mathematical Science*, 7, (1-2) (2008), 111-120.
- [13] W. M. Shah and A. Liman, On the zeros of a certain class of polynomials and related analytic functions, *Mathematica Balkanica, New Series*, 19, (2005): Facs., 3-4.